

Some properties of correlations of quantum lattice systems in thermal equilibrium

Jürg Fröhlich^{1,a)} and Daniel Ueltschi^{2,b)}

¹*Institut für Theoretische Physik, ETH Zürich, Switzerland*

²*Department of Mathematics, University of Warwick, Coventry CV4 7AL, United Kingdom*

(Received 8 December 2014; accepted 29 April 2015; published online 26 May 2015)

Simple proofs of uniqueness of the thermodynamic limit of KMS states and of the decay of equilibrium correlations are presented for a large class of quantum lattice systems at high temperatures. New quantum correlation inequalities for general Heisenberg models are described. Finally, a simplified derivation of a general result on power-law decay of correlations in 2D quantum lattice systems with continuous symmetries is given, extending results of McBryan and Spencer for the 2D classical XY model. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4921305>]

We dedicate this note to the memory of our friend Oscar E. Lanford III.

I. INTRODUCTION

Quantum lattice systems have been widely studied for many decades, heuristically, numerically, and mathematically. Many important rigorous results on equilibrium phase transitions and broken symmetries have been discovered for such systems at *low enough* temperatures. Surveys of such results can be found, e.g., in Refs. 3, 6, 8, 2, and 4 and references therein.

In this note, we study a general class of quantum lattice systems (see Sec. II) in thermal equilibrium and present simple proofs of two basic results valid at *high enough* temperatures: (i) the uniqueness of the KMS state in the thermodynamic limit and (ii) exponential decay of correlations. We also establish (iii) power-law decay of equilibrium correlations at *arbitrary* temperatures in two-dimensional quantum lattice systems with continuous symmetries. Variants of all these results have been described in the literature, see Refs. 3, 19, 16, and 22 and references therein. Our purpose, in this note, is to delineate a natural level of generality for these results and to present simple or simplified proofs thereof. Furthermore, we derive some new correlation inequalities for quantum spin systems in thermal equilibrium. These inequalities do not appear to be as useful as, e.g., the GKS- and FKG inequalities known to hold for certain classes of classical lattice systems; yet, they contain useful information on the dependence of correlations on some coupling constants. In essence, our inequalities say that correlations among spin components become stronger if the coupling constants of the interaction terms among these spin components in the Hamiltonian are increased.

II. UNIQUENESS OF KMS STATE AT HIGH TEMPERATURES

It is well-known that at sufficiently high temperatures, there are no phase-transitions, and one expects that equilibrium states are unique. This claim is backed by various mathematical results, such as analyticity of the free energy at high temperatures. In this section, we show that for a large class of quantum lattice systems, assuming that the temperature is high enough, only a single state satisfies the KMS condition (2.7) that characterizes thermal equilibrium in quantum systems. We refer the reader to the monograph of Bratteli and Robinson³ for a survey of earlier such results and references to the literature. These authors remark, in particular, that Lanford III observed that a uniqueness theorem follows from an earlier result due to Greenberg. Here, we propose to present a variant of Lanford's

^{a)}E-mail address: juerg@phys.ethz.ch

^{b)}E-mail address: daniel@ueltschi.org

approach and an improved estimate on the critical temperature. We think that our proof is somewhat simpler than the arguments described in Ref. 3. The basic idea involved in all proofs we are aware of, including ours, is to use the KMS condition to derive an inhomogeneous linear equation for the correlators of an equilibrium state satisfying the KMS condition and to show that at high enough temperatures, this equation has a unique solution (under suitable assumptions on the interactions specifying the particular quantum lattice system, see also Ref. 5).

For concreteness, we study quantum lattice systems on the simple (hyper) cubic lattice \mathbb{Z}^d . Let $\mathcal{H}_x = \mathbb{C}^N$ denote the Hilbert space of pure state vectors of the quantum-mechanical degrees of freedom, e.g., a quantum-mechanical spin, located at the site $x \in \mathbb{Z}^d$, and let $\mathcal{A}_x = \mathcal{M}_N(\mathbb{C})$ denote the algebra of bounded linear operators acting on \mathcal{H}_x , with $N < \infty$ independent of $x \in \mathbb{Z}^d$. For an arbitrary finite subset $\Lambda \subset \mathbb{Z}^d$, we define

$$\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x, \quad (2.1)$$

and we let $\mathcal{A}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{A}_x$ denote the algebra of bounded operators on \mathcal{H}_Λ . If $\Lambda \subset \Lambda'$, we view \mathcal{A}_Λ as a subalgebra of $\mathcal{A}_{\Lambda'}$ by identifying $A \in \mathcal{A}_\Lambda$ with $A \otimes \mathbb{1}_{\Lambda' \setminus \Lambda} \in \mathcal{A}_{\Lambda'}$.

Let $(\Phi_X)_{X \subset \mathbb{Z}^d}$ denote an “interaction,” that is, a collection of operators $\Phi_X \in \mathcal{A}_X$, for any finite subset X of \mathbb{Z}^d . The norm of an interaction is defined by

$$\|\Phi\|_r = \sup_{x \in \mathbb{Z}^d} \sum_{X \ni x} \|\Phi_X\| r^{|X|}. \quad (2.2)$$

Here, $\|\Phi_X\|$ denotes the usual operator norm in \mathcal{A}_X , and $r \geq 1$ is a parameter. The Hamiltonian associated with a finite domain $\Lambda \subset \mathbb{Z}^d$ is given by

$$H_\Lambda = \sum_{X \subset \Lambda} \Phi_X. \quad (2.3)$$

For $t \in \mathbb{C}$, let α_t^Λ be the linear automorphism of \mathcal{A}_Λ that describes the time evolution of operators (“observables”) in \mathcal{A}_Λ , namely,

$$\alpha_t^\Lambda(A) := e^{itH_\Lambda} A e^{-itH_\Lambda}. \quad (2.4)$$

In order to describe infinite systems, we consider the C^* -algebra, \mathcal{A} , of quasi-local observables, which is the norm-completion of the usual algebra of local observables,

$$\mathcal{A} = \overline{\mathcal{A}_0}, \quad \text{where} \quad \mathcal{A}_0 := \bigvee_{\Lambda \nearrow \mathbb{Z}^d} \mathcal{A}_\Lambda. \quad (2.5)$$

It is well-known that if $\|\Phi\|_r < \infty$, for some $r > 1$, there exists a unique one-parameter group of $*$ -automorphisms of \mathcal{A} , $\alpha_t : \mathcal{A} \rightarrow \mathcal{A}$, with $t \in \mathbb{R}$, such that

$$\lim_{n \rightarrow \infty} \|\alpha_t^{\Lambda_n}(A) - \alpha_t(A)\| = 0, \quad (2.6)$$

for an arbitrary local observable A and any sequence of domains (Λ_n) increasing to \mathbb{Z}^d , that is, such that any finite set Λ is contained in all Λ_n 's, as soon as n is large enough (depending on Λ). The operator function $\alpha_t(A)$ has an analytic continuation in t to the complex plane, for all $A \in \mathcal{A}_0$. A “state” is a bounded, positive, normalized linear functional on \mathcal{A} . A state ρ describes *thermal equilibrium at inverse temperature β* iff it satisfies the KMS condition, i.e., iff

$$\rho(AB) = \rho(B \alpha_{i\beta}(A)), \quad (2.7)$$

for all A, B in \mathcal{A}_0 . By considering sequences of finite-volume (Gibbs) equilibrium states, a standard compactness argument shows the existence of cluster points of states that satisfy the KMS condition, i.e., the existence of KMS states is an almost trivial fact. We are now prepared to state our uniqueness theorem.

Theorem 2.1. *Assume that*

$$\beta \|\Phi\|_{N+1} < (2N)^{-1}.$$

Then, there exists a unique KMS state at inverse temperature β .

We actually prove the theorem under the more general condition that there exists $s < 1/N$ such that $2\beta\|\Phi\|_{N(1+s)} < s$. As mentioned above, the strategy of our proof is to reformulate the KMS condition as an equation for the equilibrium state that has a unique solution when β is small enough. In order to derive this equation, we express observables as commutators of operators. The proof of Theorem 2.1 will be given after the one of Lemma 2.2, which we state next.

Here and in the sequel, $\|\cdot\|_{\text{HS}}$ denotes the normalized Hilbert-Schmidt norm,

$$\|A\|_{\text{HS}}^2 = \frac{1}{\dim \mathcal{H}_\Lambda} \text{Tr } A^*A. \tag{2.8}$$

Notice that

$$\frac{1}{\sqrt{\dim \mathcal{H}_\Lambda}} \|A\| \leq \|A\|_{\text{HS}} \leq \|A\| \tag{2.9}$$

for all $A \in \mathcal{A}_\Lambda$.

Lemma 2.2. *Let A be a hermitian $N \times N$ matrix with the property that $\text{Tr } A = 0$. Then, there exist hermitian $N \times N$ matrices B_1, \dots, B_{N-1} and C_1, \dots, C_{N-1} such that*

$$A = \sum_{i=1}^{N-1} [B_i, C_i],$$

$$\sum_{i=1}^{N-1} \|B_i\|_{\text{HS}} \|C_i\|_{\text{HS}} \leq \sqrt{N} \|A\|_{\text{HS}}.$$

Proof. Let a_1, \dots, a_N be the eigenvalues of A (repeated according to their multiplicity). We have that

$$\sum_{i=1}^N a_i = 0, \quad \sum_{i=1}^N |a_i|^2 = N \|A\|_{\text{HS}}^2. \tag{2.10}$$

In particular, each $|a_i|$ is bounded above by $\sqrt{N}\|A\|_{\text{HS}}$. Let us order the eigenvalues so that

$$\left| \sum_{i=1}^k a_i \right| \leq \sqrt{N} \|A\|_{\text{HS}} \tag{2.11}$$

for all $1 \leq k \leq N - 1$. This is indeed possible, as can be seen by induction using $\sum a_i = 0$: if $0 \leq \sum_{i=1}^k a_i \leq \sqrt{N}\|A\|_{\text{HS}}$, we can find $a_{k+1} \leq 0$ among the remaining eigenvalues such that $|\sum_{i=1}^{k+1} a_i| \leq \sqrt{N}\|A\|_{\text{HS}}$. And if the partial sum is negative, we can find $a_{k+1} \geq 0$ among the remaining eigenvalues, with the same conclusion.

We work in a basis such that A is diagonal and its eigenvalues are ordered so they satisfy the properties above. Let $\tilde{a}_k = \sum_{i=1}^k a_i$, and let $\sigma_{j,j+1}^1, \sigma_{j,j+1}^2, \sigma_{j,j+1}^3$ be $N \times N$ matrices that are equal to Pauli matrices on the 2×2 block that contains (j, j) and $(j + 1, j + 1)$ and that are equal to zero everywhere else. It is not hard to check that

$$A = \sum_{j=1}^{N-1} \tilde{a}_j \sigma_{j,j+1}^3. \tag{2.12}$$

We therefore have that

$$A = \frac{1}{2} \sum_{j=1}^{N-1} \tilde{a}_j [\sigma_{j,j+1}^1, \sigma_{j,j+1}^2], \tag{2.13}$$

which proves the first claim. The bound follows from $|\tilde{a}_j| \leq \sqrt{N}\|A\|_{\text{HS}}$ and $\|\sigma_{j,j+1}^i\|_{\text{HS}}^2 = 2/N$. \square

Proof of Theorem 2.1. Let $(e_i)_{i=0}^{N^2-1}$ be a hermitian basis of $\mathcal{M}_N(\mathbb{C})$, with $e_0 = \mathbb{1}$, $\text{Tr } e_i = 0$ if $1 \neq 0$, and $\|e_i\| = 1$, for all i . Let J be the set of multi-indices $j = (j_x)_{x \in \mathbb{Z}^d}$, $0 \leq j_x \leq N^2 - 1$, with

finite support

$$\text{supp } j = \{x \in \mathbb{Z}^d | j_x \neq 0\}. \tag{2.14}$$

Given $j \in J$, let $e_j = \otimes_{x \in \text{supp } j} e_{j_x} \in \mathcal{A}_{\text{supp } j}$. The linear span of $\{e_j\}_{j \in J}$ is dense in \mathcal{A} .

Let tr denote the normalized trace on \mathcal{A} ; it is equal to $\frac{1}{\dim \mathcal{H}_\Lambda}$ Tr on \mathcal{A}_Λ and it can be extended to \mathcal{A} by continuity. The state ρ can be written as $\rho = \text{tr} + \varepsilon$, where $\varepsilon(\mathbb{1}) = 0$. We actually have that

$$\varepsilon(e_j) = \begin{cases} \rho(e_j) & \text{if } j \neq 0, \\ 0 & \text{if } j \equiv 0. \end{cases} \tag{2.15}$$

Using Lemma 2.2, we have that

$$e_j = \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} [\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)}, \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}], \tag{2.16}$$

for $j \neq 0$. Here, $b_i^{(k)}, c_i^{(k)}$ are the matrices B_i, C_i of Lemma 2.2 in the case where the matrix A is e_k .

We now use this decomposition and KMS condition (2.7) in order to get an equation for ε . For $j \neq 0$,

$$\begin{aligned} \varepsilon(e_j) &= \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} \rho([\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)}, \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}]) \\ &= \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} \rho(\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)} \cdot (\mathbb{1} - \alpha_{i\beta}) \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}) \\ &= \delta(e_j) + K_\beta \varepsilon(e_j). \end{aligned} \tag{2.17}$$

In the above equation, we set

$$\delta(e_j) = \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} \text{tr} (\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)} \cdot (\mathbb{1} - \alpha_{i\beta}) \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}), \tag{2.18}$$

and the operator K_β is defined by

$$(K_\beta \phi)(e_j) = \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} \phi(\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)} \cdot (\mathbb{1} - \alpha_{i\beta}) \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}). \tag{2.19}$$

Notice that K_β is a linear operator on the Banach space $\mathcal{L}(\mathcal{A})$ of linear functionals on \mathcal{A} . Equation (2.17) can be written as

$$(\mathbb{1} - K_\beta)\varepsilon = \delta. \tag{2.20}$$

Let us introduce the following norm on $\mathcal{L}(\mathcal{A})$:

$$\|\phi\| = \sup_{j \in J} |\phi(e_j)|. \tag{2.21}$$

Because $\|e_j\| = 1$ for all j , we have $\|\phi\| \leq \|\phi\|$ and $(\mathcal{L}(\mathcal{A}), \|\cdot\|)$ is a normed vector space. We consider K_β as an operator on $(\mathcal{L}(\mathcal{A}), \|\cdot\|)$ and we show that its norm is strictly less than 1; the solution of (2.20) is then unique. The norm of K_β is equal to

$$\|K_\beta\| = \sup_{\|\phi\|=1} \sup_{j \in J} |K_\beta \phi(e_j)|. \tag{2.22}$$

Recall that $\alpha_{i\beta} = \lim_\Lambda \alpha_{i\beta}^\Lambda$ (with convergence in the operator norm) and that $\alpha_{i\beta}^\Lambda(A)$, $A \in \mathcal{A}$, has a well-known expansion in multiple commutators. From (2.19), we get

$$\begin{aligned} |K_\beta \phi(e_j)| \leq & \frac{1}{|\text{supp } j|} \sum_{y \in \text{supp } j} \sum_{i=1}^{N-1} \sum_{n \geq 1} \frac{\beta^n}{n!} \sup_{\Lambda \subset \mathbb{Z}^d} \sum_{X_1, \dots, X_n \subset \Lambda} \\ & \left| \phi(\otimes_{x \neq y} e_{j_x} \otimes b_i^{(jy)} [\Phi_{X_n}, \dots, [\Phi_{X_1}, \otimes_{x \neq y} \mathbb{1} \otimes c_i^{(jy)}] \dots]) \right|. \end{aligned} \tag{2.23}$$

Because of the commutators, the sum over the X_k 's is restricted to subsets that satisfy

$$\begin{aligned} X_1 \ni y, \\ X_2 \cap X_1 \neq \emptyset, \\ \vdots \\ X_n \cap (X_1 \cup \dots \cup X_{n-1}) \neq \emptyset. \end{aligned} \tag{2.24}$$

Let $A = \sum_{(j'_x)_{x \in X}} a_{j'_x} e_{j'_x}$ be an operator in \mathcal{A}_X . For any $(j_x)_{x \in X}$, we have

$$\begin{aligned} |\phi(\otimes_{x \in X} e_{j_x} \otimes A)| &= \left| \sum_{(j'_x)_{x \in X}} a_{j'_x} \phi(\otimes_{x \in X} e_{j_x} \otimes e_{j'_x}) \right| \\ &\leq \|\phi\| \sum_{(j'_x)_{x \in X}} |a_{j'_x}| \\ &\leq \|\phi\| \|A\|_{\text{HS}} N^{|X|}. \end{aligned} \tag{2.25}$$

Using Eq. (2.25) with $\|\phi\| = 1$, $\|AB\|_{\text{HS}} \leq \|A\| \|B\|_{\text{HS}}$, and $\|c_i^{(j_y)}\| \leq \sqrt{N} \|c_i^{(j_y)}\|_{\text{HS}}$, we get

$$\begin{aligned} |K_\beta \phi(e_j)| &\leq \sqrt{N} \sup_{y \in \mathbb{Z}^d} \sum_{n \geq 1} \frac{(2\beta)^n}{n!} \sum_{X_1, \dots, X_n: y} \left(\prod_{k=1}^n \|\Phi_{X_k}\| N^{|X_k|} \right) \sum_{i=1}^{N-1} \|b_i^{(j_y)}\|_{\text{HS}} \|c_i^{(j_y)}\|_{\text{HS}} \\ &\leq N \sup_{y \in \mathbb{Z}^d} \sum_{n \geq 1} \frac{(2\beta)^n}{n!} \sum_{X_1, \dots, X_n: y} \prod_{k=1}^n \|\Phi_{X_k}\| N^{|X_k|}. \end{aligned} \tag{2.26}$$

We have used Lemma 2.2 to get the last line. The constraint $X_1, \dots, X_n : y$ means that (2.24) must be respected. The final step is to estimate the sum over such subsets. This can be conveniently done with an inductive argument. Namely, let $R_0 = 0$ and, for $m \geq 1$, let

$$R_m = \sup_{y \in \mathbb{Z}^d} \sum_{n=1}^m \frac{(2\beta)^n}{n!} \sum_{X_1, \dots, X_n: y} \prod_{k=1}^n \|\Phi_{X_k}\| N^{|X_k|}. \tag{2.27}$$

Summing first over $X_1 \ni y$, then over sets that intersect sites of X_1 , we get

$$\begin{aligned} R_m &\leq 2\beta \sup_y \sum_{X_1 \ni y} \|\Phi_{X_1}\| N^{|X_1|} \prod_{x \in X_1} \left(\sum_{n=1}^m \frac{(2\beta)^{n-1}}{(n-1)!} \sum_{X_2, \dots, X_n: x} \prod_{k=2}^n \|\Phi_{X_k}\| N^{|X_k|} \right) \\ &\leq 2\beta \sup_y \sum_{X_1 \ni y} \|\Phi_{X_1}\| N^{|X_1|} (1 + R_{m-1})^{|X_1|}. \end{aligned} \tag{2.28}$$

It follows easily that $R_m \leq r$ for all m and all r such that $2\beta \|\Phi\|_{N(1+r)} \leq r$. Then, $\|K_\beta\| \leq Nr$, and the assumption of Theorem 2.1 implies the existence of r such that $Nr < 1$.

III. HIGH TEMPERATURE EXPANSIONS

(Connected) correlations between operators localized in disjoint regions of the lattice vanish when $\beta = 0$. For positive, but small β and short-range interactions, correlations decay exponentially fast. This can be proven in several different ways. Here, we use the method of *cluster expansions*, which is robust and applies to both classical and quantum systems. The main result of this section and our method of proof are not new; see Ref. 21, Sec. V.5, and references therein. Our approach is based on the simple exposition in Ref. 22. It is quite direct and straightforward.

As an alternative to cluster expansions, one should mention the method of Lee and Yang, i.e., general *Lee-Yang theorems*. This method establishes and then exploits analyticity properties of correlation functions in variables corresponding to external magnetic fields. It yields exponential decay of correlations, provided the magnetic field variables belong to certain subsets of the complex plane. We do not wish to describe these matters in more detail here, but see Refs. 18, 13, 20, and 11 for precise statements of results and proofs.

A. Analyticity of the free energy

Let Λ be a finite subset of \mathbb{Z}^d . Let \mathcal{S}_Λ denote the set of finite sequences (X_1, \dots, X_n) , with $n \geq 1$ and $X_i \subset \Lambda$ for all i . Let $\mathcal{C}_\Lambda \subset \mathcal{S}_\Lambda$ denote the set of *clusters*, i.e., the set of objects $C = (X_1, \dots, X_n)$ such that the graph

$$\{(i, j) : X_i \cap X_j \neq \emptyset\} \tag{3.1}$$

is connected. We also let $\text{supp } C = \cup_i X_i$ denote the support of C . We introduce the following weight function on \mathcal{S}_Λ : if $C = (X_1, \dots, X_n)$,

$$w(C) = \frac{\beta^n}{n!} \text{tr } \Phi_{X_1} \dots \Phi_{X_n}. \tag{3.2}$$

Finally, let φ denote the usual combinatorial function of cluster expansions, namely,

$$\varphi(C_1, \dots, C_k) = \begin{cases} 1 & \text{if } k = 1, \\ \sum_{g \in \text{Conn}(k)} \prod_{\{i,j\} \in g} (-1)^{\text{supp } C_i \cap \text{supp } C_j \neq \emptyset} & \text{if } k \geq 2. \end{cases} \tag{3.3}$$

Here, $\text{Conn}(k)$ is the set of connected graphs of k vertices, and the product is over the edges of the connected graph g .

The first result deals with the partition function

$$Z_\Lambda = \text{tr } e^{-\beta H_\Lambda}, \tag{3.4}$$

with H_Λ the Hamiltonian defined in Eq. (2.3). As before, tr denotes the normalized trace. It follows easily from Theorem 3.1 that the free energy $f_\Lambda(\beta) = -\frac{1}{|\Lambda|} \log Z_\Lambda$ is analytic in β in the infinite-volume limit.

Theorem 3.1. *Assume that there exists $a > 0$ such that*

$$\beta \|\Phi\| e^{a\beta} \leq a.$$

Then, the partition function has the expression

$$Z_\Lambda = \exp \left\{ \sum_{k \geq 1} \frac{1}{k!} \sum_{C_1, \dots, C_k \in \mathcal{C}_\Lambda} \varphi(C_1, \dots, C_k) \prod_{i=1}^k w(C_i) \right\}.$$

The sums are absolutely convergent, and

$$1 + \sum_{k \geq 2} k \sum_{C_1, \dots, C_k \in \mathcal{C}_\Lambda} |\varphi(C_2, \dots, C_k)| \prod_{i=2}^k |w(C_i)| \leq e^{a|\text{supp } C_1|},$$

for all $\Lambda \subset \mathbb{Z}^d$ and all $C_1 \in \mathcal{C}_\Lambda$.

We remark that, historically, the ‘‘clusters’’ of the expansion are the connected sets of $\text{Conn}(k)$ in Eq. (3.3) rather than our C_i ’s. Clusters are often grouped according to their supports, which yields the ‘‘polymer’’ expansion. But we find it better to keep the C_i ’s as they are, without resummation.

Proof. Clearly,

$$Z_\Lambda = \text{tr } e^{-\beta \sum_{X \subset \Lambda} \Phi_X} = \sum_{n \geq 0} \frac{\beta^n}{n!} \sum_{X_1, \dots, X_n \subset \Lambda} \text{tr } \Phi_{X_1} \dots \Phi_{X_n}. \tag{3.5}$$

We group the sets X_1, \dots, X_n in clusters. We get

$$Z_\Lambda = \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in \mathcal{C}_\Lambda \\ \text{disjoint}}} w(C_1) \dots w(C_k). \tag{3.6}$$

The sum is restricted on “disjoint” clusters such that $\text{supp } C_i \cap \text{supp } C_j = \emptyset$ for all $i \neq j$. This expression fits the framework of the method of cluster expansion. A sufficient condition for its convergence^{12,16,22} is that there exists $a > 0$ such that

$$\sum_{\substack{C' \in \mathcal{C}_\Lambda \\ \text{supp } C' \cap \text{supp } C \neq \emptyset}} |w(C')| e^{a|\text{supp } C'} \leq a|\text{supp } C|, \tag{3.7}$$

for all $C \in \mathcal{C}_\Lambda$. Once (3.7) is proved, Theorem 3.1 follows immediately from e.g. Ref. 22, Theorem 1.

Let $n(C)$ denote the number of sets that constitute the cluster C . We have

$$|w(C)| \leq \frac{\beta^{n(C)}}{n(C)!} \prod_{i=1}^{n(C)} \|\Phi_{X_i}\|. \tag{3.8}$$

Let $R_0 = 0$, and, for $m \geq 1$,

$$R_m = \sup_{x \in \mathbb{Z}^d} \sum_{\substack{C \in \mathcal{C}_\Lambda, \text{supp } C \ni x \\ n(C) \leq m}} \frac{\beta^{n(C)}}{n(C)!} \prod_{i=1}^{n(C)} \|\Phi_{X_i}\| e^{a|\text{supp } C}|. \tag{3.9}$$

We show that $R_m \leq a$ for all m (and all Λ); this implies (3.7). We prove it by induction by means of inequality (3.11) below. We now give a careful derivation.

Let $x \in \Lambda$, and let us consider an order on the subsets of Λ with the property that $X < X'$ if $X \ni x \notin X'$. If f is a non-negative function on subsets of Λ , and writing $f(C) = \prod f(X_i)$, we have

$$\begin{aligned} \frac{1}{n!} \sum_{\substack{C=(X_1, \dots, X_n) \in \mathcal{C}_\Lambda \\ \text{supp } C \ni x}} \prod_{i=1}^n f(X_i) &\leq \sum_{\substack{X_1 < \dots < X_n \\ X_1 \ni x, (X_1, \dots, X_n) \in \mathcal{C}_\Lambda}} \prod_{i=1}^n f(X_i) \\ &= \sum_{X_1 \ni x} f(X_1) \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in \mathcal{C}_\Lambda, \text{disjoint} \\ n(C_1) + \dots + n(C_k) = n-1 \\ X_1 < C_i, \text{supp } C_i \cap X_1 = \emptyset \forall i}} \prod_{i=1}^k \frac{1}{n(C_i)!} f(C_i). \end{aligned} \tag{3.10}$$

The inequality in the first line is due to the case of identical sets, $X_i = X_j$ for some $i \neq j$. In the last sum, the constraint $X_1 < C_i$ means that X_1 is smaller than all the sets of C_i . It follows that

$$\begin{aligned} R_m &\leq \beta \sup_{x \in \mathbb{Z}^d} \sum_{X \ni x} \|\Phi_X\| e^{a|X|} \prod_{y \in X} \left(1 + \sum_{\substack{C \in \mathcal{C}_\Lambda, \text{supp } C \ni y \\ n(C) \leq m-1}} \frac{\beta^{n(C)}}{n(C)!} \prod_{i=1}^{n(C)} \|\Phi_{X_i}\| e^{a|\text{supp } C}| \right) \\ &\leq \beta \sup_{x \in \mathbb{Z}^d} \sum_{X \ni x} \|\Phi_X\| (e^a (1 + R_{m-1}))^{|X|}. \end{aligned} \tag{3.11}$$

Using the induction hypothesis $R_{m-1} \leq a$ and the assumption of the theorem, we get $R_m \leq a$. This proves (3.7). □

B. Thermodynamic limit and expectations of local observables

Next, we consider the expectation of observables. Let $A \in \mathcal{A}_\Lambda$. We let $\text{supp } A$ denote the support of the observable A ; it is equal to the smallest set X such that $A \in \mathcal{A}_X$. We are interested in the expectation

$$\langle A \rangle = \frac{1}{Z_\Lambda} \text{tr } A e^{-\beta H_\Lambda}. \tag{3.12}$$

A similar expansion than (3.5) gives

$$\text{tr } A e^{-\beta H_\Lambda} = \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_A, C_1, \dots, C_k \\ \text{disjoint}}} w_A(C_A) w(C_1) \dots w(C_k), \tag{3.13}$$

where

$$w_A(C_A) = \frac{\beta^n}{n!} \text{tr } A\Phi_{X_1} \dots \Phi_{X_n}. \tag{3.14}$$

Here, $C_A = (X_0, X_1, \dots, X_n)$ is a cluster such that $X_0 = \text{supp } A$ by definition. $n = 0$ is possible in which case $w_A(C_A) = \text{tr } A$.

Under the same assumption as in Theorem 3.1, the method of cluster expansion applies and it gives

$$\text{tr } A e^{-\beta H_\Lambda} = \sum_{C_A} w_A(C_A) \exp \left\{ \sum_{k \geq 1} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in C_\Lambda \\ C_A, C_1, \dots, C_k \text{ disjoint}}} \varphi(C_1, \dots, C_k) \prod_{i=1}^k w(C_i) \right\}. \tag{3.15}$$

This can be combined with the expression for Z_Λ in Theorem 3.1, because of cancellations, we obtain

$$\langle A \rangle = \sum_{C_A} w_A(C_A) \exp \left\{ \sum_{k \geq 1} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in C_\Lambda \\ (\cup_i \text{supp } C_i) \cap \text{supp } C_A \neq \emptyset}} \varphi(C_1, \dots, C_k) \prod_{i=1}^k w(C_i) \right\}. \tag{3.16}$$

This expression makes it possible to take the thermodynamic limit $\Lambda \nearrow \mathbb{Z}^d$ as all sums converge absolutely and uniformly.

C. Decay of two-point correlation functions

Let $b(X)$ be a non-negative function of finite subsets of \mathbb{Z}^d . The larger this function, the better the decay. We assume a slightly stronger condition on the interaction, namely, that there exists $a > 0$ such that

$$\beta \sup_{x \in \mathbb{Z}^d} \sum_{X \ni x} \|\Phi_X\| e^{\frac{3}{2}a|X| + b(X)} < a. \tag{3.17}$$

Given two sets $X, Y \subset \mathbb{Z}^d$, let

$$\mu(X, Y) = \min_{n \geq 1} \min_{\substack{X_1, \dots, X_n \\ X_i \cap X_{i+1} \neq \emptyset \forall i=0, \dots, n+1}} b(X_1) + \dots + b(X_n). \tag{3.18}$$

In the second minimum, we set $X_0 = X$ and $X_{n+1} = Y$.

Theorem 3.2. *Assume that the interaction Φ satisfies condition (3.17). Then, we have*

$$|\langle AB \rangle - \langle A \rangle \langle B \rangle| \leq k(A, B) e^{-\mu(\text{supp } A, \text{supp } B)},$$

with

$$k(A, B) = \|A\| \|B\| (a|\text{supp } A| + a|\text{supp } B| + 3a^2|\text{supp } A| |\text{supp } B|).$$

As A and B are moved away from each other, the decay is given by $e^{-\mu(\cdot)}$. Decay is exponential if the interactions are finite-range or exponentially decaying.

Proof. An expansion similar to (3.13) holds in the case where A is replaced by the product of two operators, AB . We denote C_{AB} the clusters of the type $(\text{supp } A, \text{supp } B, X_1, \dots, X_n)$; $n = 0$ is not possible unless $\text{supp } A \cap \text{supp } B \neq \emptyset$. The corresponding weight is

$$w_{AB}(C_{AB}) = \frac{\beta^n}{n!} \text{tr } AB\Phi_{X_1} \dots \Phi_{X_n}. \tag{3.19}$$

Expansion of the exponential gives

$$\begin{aligned} \text{tr } AB e^{-\beta H_\Lambda} &= \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_{AB}, C_1, \dots, C_k \\ \text{disjoint}}} w_{AB}(C_{AB}) w(C_1) \dots w(C_k) \\ &+ \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_A, C_B, C_1, \dots, C_k \\ \text{disjoint}}} \frac{1}{k!} w_A(C_A) w_B(C_B) w(C_1) \dots w(C_k). \end{aligned} \tag{3.20}$$

It is convenient to use the following notation, which mirrors that of Ref. 22, Sec. 3:

$$\begin{aligned} Z_\Lambda(C_A) &= \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in C_\Lambda \setminus \text{supp } C_A \\ \text{disjoint}}} w(C_1) \dots w(C_k), \\ Z_\Lambda(C_A, C_B) &= \sum_{k \geq 0} \frac{1}{k!} \sum_{\substack{C_1, \dots, C_k \in C_\Lambda \setminus (\text{supp } C_A \cup \text{supp } C_B) \\ \text{disjoint}}} w(C_1) \dots w(C_k), \\ \hat{Z}_\Lambda(C_A) &= \sum_{k \geq 0} (k+1) \sum_{C_1, \dots, C_k \in C_\Lambda} \varphi(C_A, C_1, \dots, C_k) w(C_1) \dots w(C_k), \\ \hat{Z}_\Lambda(C_A, C_B) &= \sum_{k \geq 0} (k+1)(k+2) \sum_{C_1, \dots, C_k \in C_\Lambda} \varphi(C_A, C_B, C_1, \dots, C_k) w(C_1) \dots w(C_k). \end{aligned} \tag{3.21}$$

We have

$$\text{tr } AB e^{-\beta H_\Lambda} = \sum_{C_{AB}} w_{AB}(C_{AB}) Z_\Lambda(C_{AB}) + \sum_{\substack{C_A, C_B \\ \text{disjoint}}} w_A(C_A) w_B(C_B) Z_\Lambda(C_A, C_B). \tag{3.22}$$

It follows from Ref. 22, Theorem 2, that

$$\begin{aligned} \langle AB \rangle - \langle A \rangle \langle B \rangle &= \sum_{C_{AB}} w_{AB}(C_{AB}) \hat{Z}_\Lambda(C_{AB}) + \sum_{\substack{C_A, C_B \\ \text{disjoint}}} w_A(C_A) w_B(C_B) \hat{Z}_\Lambda(C_A, C_B) \\ &- \sum_{\substack{C_A, C_B \\ \text{supp } C_A \cap \text{supp } C_B \neq \emptyset}} w_A(C_A) w_B(C_B) \hat{Z}_\Lambda(C_A) \hat{Z}_\Lambda(C_B). \end{aligned} \tag{3.23}$$

Let $b(C) = \sum_i b(X_i)$ for $C = (X_1, \dots, X_n)$. Adapting the proof of (3.7), one can show that

$$\sum_{\substack{C' \in C_\Lambda \\ \text{supp } C' \cap \text{supp } C \neq \emptyset}} |w(C')| e^{\frac{3}{2}a|\text{supp } C'| + b(C')} \leq a|\text{supp } C|. \tag{3.24}$$

This allows to use Ref. 22, Theorem 3. For $C_A = (\text{supp } A, X_1, \dots, X_n)$, we get

$$e^{b(C_A)} |\hat{Z}_\Lambda(C_A)| \leq e^{a|\text{supp } C_A| + b(\text{supp } A)}. \tag{3.25}$$

The same bound applies to C_B ; as for C_{AB} , we have

$$e^{\mu(\text{supp } C_A, \text{supp } C_B)} |\hat{Z}_\Lambda(C_A, C_B)| \leq e^{\frac{3}{2}a|\text{supp } C_A| + \frac{3}{2}a|\text{supp } C_B|}. \tag{3.26}$$

Theorem 3.2 follows from expression (3.23) and bounds (3.24)–(3.26). □

IV. CORRELATION INEQUALITIES FOR QUANTUM SPIN SYSTEMS

We now consider a more restricted setting. Let S^1, S^2, S^3 be spin operators in \mathbb{C}^N that satisfy $[S^1, S^2] = iS^3$ and with the other commutation relations obtained by cyclic permutations of indices. Let $S_x^i = S^i \otimes \text{Id}_{\Lambda \setminus \{x\}}$, $i = 1, 2, 3$. The Hamiltonian depends on real coupling parameters (exchange couplings), $J_{xy}^i = J_{yx}^i$, and is given by

$$H_\Lambda = -\frac{1}{2} \sum_{x, y \in \Lambda} (J_{xy}^1 S_x^1 S_y^1 + J_{xy}^2 S_x^2 S_y^2 + J_{xy}^3 S_x^3 S_y^3). \tag{4.1}$$

Here, Λ is an arbitrary finite set. From now on, we use the usual trace, denoted Tr , rather than the normalized trace, tr . With $Z_\Lambda = \text{Tr} e^{-\beta H_\Lambda}$ denoting the partition function, the correlation functions at inverse temperature β are given by

$$\langle S_0^i S_x^i \rangle = \frac{1}{Z_\Lambda} \text{Tr} (S_0^i S_x^i e^{-\beta H_\Lambda}). \tag{4.2}$$

The case $J_{xy}^1 = J_{xy}^2 = 0$, for all $x, y \in \Lambda$, corresponds to the Ising model, which is in fact a classical model. The case $J_{xy}^3 = 0$ and $J_{xy}^1 = J_{xy}^2$, for all x, y , corresponds to the quantum XY model. And the symmetric case, $J_{xy}^1 = J_{xy}^2 = J_{xy}^3$, corresponds to the isotropic Heisenberg model. Positive values of the couplings correspond to ferromagnetic order, while negative values of the couplings correspond to antiferromagnetism.

It is natural to expect that correlations are stronger among those components of the spins that correspond to stronger coupling parameters in the Hamiltonian. This is the content of the next theorem. The inequalities stated there do not appear to have been noticed before, except for the spin- $\frac{1}{2}$ XY model corresponding to $N = 2$: assuming that Λ is a rectangular subset of \mathbb{Z}^d and $\|x\| = 1$, the first inequality follows from reflection positivity;¹⁷ for general Λ and general x , it follows from a random loop representation.²³

Theorem 4.1. *Assume that, for all $x, y \in \Lambda$, the couplings satisfy*

$$|J_{xy}^2| \leq J_{xy}^1.$$

Then, we have that

$$|\langle S_0^2 S_x^2 \rangle| \leq \langle S_0^1 S_x^1 \rangle,$$

for all $x \in \Lambda$. More generally, for all $x_1, \dots, x_k \in \Lambda$ and $j_1, \dots, j_k \in \{1, 2\}$,

$$|\langle S_{x_1}^{j_1} \dots S_{x_k}^{j_k} \rangle| \leq \langle S_{x_1}^1 \dots S_{x_k}^1 \rangle.$$

Further inequalities can be generated using symmetries. Some inequalities hold for the staggered two-point function $(-1)^{|x|} \langle S_0^i S_x^i \rangle$.

Proof. Let $S \in \frac{1}{2}\mathbb{N}$ such that $2S + 1 = N$, and let $|a\rangle, a \in \{-S, \dots, S\}$ denote basis elements of \mathbb{C}^{2S+1} . Let the operators S^\pm be defined by

$$S^+|a\rangle = \sqrt{S(S+1) - a(a+1)} |a+1\rangle, \quad S^-|a\rangle = \sqrt{S(S+1) - (a-1)a} |a-1\rangle, \tag{4.3}$$

with the understanding that $S^+|S\rangle = S^-| -S\rangle = 0$. Then, let $S^1 = \frac{1}{2}(S^+ + S^-)$, $S^2 = \frac{1}{2i}(S^+ - S^-)$, and $S^3|a\rangle = a|a\rangle$. It is well-known that these operators satisfy the spin commutation relations. Further, the matrix elements of S^1, S^\pm are all non-negative, and the matrix elements of S^2 are all less than or equal to those of S^1 in absolute values. Using the Trotter formula and multiple resolutions of the identity, we have

$$\begin{aligned} |\text{Tr} S_0^2 S_x^2 e^{-\beta H_\Lambda}| &\leq \lim_{N \rightarrow \infty} \sum_{\sigma_0, \dots, \sigma_N \in \{-S, \dots, S\}^\Lambda} |\langle \sigma_0 | S_0^2 S_x^2 | \sigma_1 \rangle \\ &\quad \langle \sigma_1 | e^{\frac{\beta}{N} \sum J_{yz}^3 S_y^3 S_z^3} | \sigma_1 \rangle \langle \sigma_1 | \left(1 + \frac{\beta}{N} \sum_{y,z \in \Lambda} (J_{yz}^1 S_y^1 S_z^1 + J_{yz}^2 S_y^2 S_z^2) \right) | \sigma_2 \rangle \\ &\quad \dots \langle \sigma_N | e^{\frac{\beta}{N} \sum J_{yz}^3 S_y^3 S_z^3} | \sigma_N \rangle \langle \sigma_N | \left(1 + \frac{\beta}{N} \sum_{y,z \in \Lambda} (J_{yz}^1 S_y^1 S_z^1 + J_{yz}^2 S_y^2 S_z^2) \right) | \sigma_0 \rangle |. \end{aligned} \tag{4.4}$$

Observe that the matrix elements of all operators are non-negative, except for $S_0^2 S_x^2$. Indeed, this follows from

$$J_{yz}^1 S_y^1 S_z^1 + J_{yz}^2 S_y^2 S_z^2 = \frac{1}{4} (J_{yz}^1 - J_{yz}^2) (S_y^+ S_z^+ + S_y^- S_z^-) + \frac{1}{4} (J_{yz}^1 + J_{yz}^2) (S_y^+ S_z^- + S_y^- S_z^+). \tag{4.5}$$

We get an upper bound for the right side of (4.4) by replacing $|\langle \sigma_0 | S_0^2 S_x^2 | \sigma_1 \rangle|$ with $\langle \sigma_0 | S_0^1 S_x^1 | \sigma_1 \rangle$. We have obtained

$$|\mathrm{Tr} S_0^2 S_x^2 e^{-\beta H_\Lambda}| \leq \mathrm{Tr} S_0^1 S_x^1 e^{-\beta H_\Lambda}, \quad (4.6)$$

which proves the first claim. The second claim can be proved exactly the same way. \square

Corollary 4.2. Assume that for all $x, y \in \Lambda$, the couplings satisfy

$$J_{xy}^1 = J_{xy}^2 \geq 0.$$

Then, we have for all $x, y, z, u \in \Lambda$,

$$\frac{\partial}{\partial J_{xy}^1} \langle S_z^2 S_u^2 \rangle \leq \frac{\partial}{\partial J_{xy}^1} \langle S_z^1 S_u^1 \rangle.$$

Proof. For $i = 1, 2, 3$, we have

$$\frac{1}{\beta} \frac{\partial}{\partial J_{xy}^1} \langle S_z^i S_u^i \rangle = (S_x^1 S_y^1, S_z^i S_u^i) - \langle S_x^1 S_y^1 \rangle \langle S_z^i S_u^i \rangle, \quad (4.7)$$

where (A, B) denotes the Duhamel two-point function,

$$(A, B) = \frac{1}{Z_\Lambda} \int_0^1 \mathrm{Tr} A e^{-s\beta H_\Lambda} B e^{-(1-s)\beta H_\Lambda} ds. \quad (4.8)$$

It is not hard to extend the proof of Theorem 4.1 to the Duhamel function, so that

$$|(S_x^1 S_y^1, S_z^2 S_u^2)| \leq (S_x^1 S_y^1, S_z^1 S_u^1). \quad (4.9)$$

Further, we have $\langle S_z^2 S_u^2 \rangle = \langle S_z^1 S_u^1 \rangle$ by symmetry. The result follows. \square

V. DECAY OF CORRELATIONS DUE TO SYMMETRIES

In this section, we prove a variant of the Mermin-Wagner theorem. Our method of proof only works for systems that are effectively two-dimensional. The first result, with an explicit bound on the two-point correlation function, is due to Fisher and Jasnow.⁷ Unfortunately, it only yields logarithmic decay. The decay is, however, expected to be power-law, and this was proven by McBryan and Spencer¹⁹ in a short and lucid article that exploits complex rotations. Power-law decay was proven for some quantum systems in Refs. 1 and 14. The proofs use Fourier transform and the Bogoliubov inequality, and they are limited to regular two-dimensional lattices. A much more general result was obtained by Koma and Tasaki using complex rotations.¹⁵ The present proof is similar to theirs but somewhat simpler. Absence of ordering and of symmetry breaking was proven in Refs. 9 and 10.

We assume that $J_{xy}^1 = J_{xy}^2$ for all x, y . The decay of correlations is measured by the following expression:

$$\xi_\beta(x) = \sup_{\substack{(\phi_y) \in \mathbb{R}^\Lambda \\ \phi_x = 0}} \left[\phi_0 - 2\beta S^2 \sum_{y, z \in \Lambda} |J_{yz}^1| (\cosh(\phi_y - \phi_z) - 1) \right]. \quad (5.1)$$

The solution of this variational problem is essentially a discrete harmonic function. We can estimate it explicitly in the case of “2D-like” graphs with nearest-neighbor couplings. Let Λ denote a graph, i.e., a finite set of vertices and a set of edges, and let $d(x, y)$ denote the graph distance, i.e., the length of the shortest path that connects x and y .

Lemma 5.1. Assume that $J_{xy}^i = 0$ whenever $d(x, y) \neq 1$ and let $J = \max |J_{xy}^i|$. Assume in addition that there exists a constant K such that, for any $\ell \geq 0$,

$$\#\{\{x, y\} \subset \Lambda : d(0, x) = \ell, d(0, y) = \ell + 1, \text{ and } d(x, y) = 1\} \leq K(\ell + 1).$$

Then, there exists $C = C(\beta, S, J, K)$, which does not depend on x , such that

$$\xi_\beta(x) \geq \frac{1}{16\beta JS^2 K} \log(d(0, x) + 1) - C.$$

Proof. With c to be chosen later, let

$$\phi_y = \begin{cases} c \log \frac{d(0,x)+1}{d(0,y)+1} & \text{if } d(0,y) \leq d(0,x), \\ 0 & \text{otherwise.} \end{cases} \tag{5.2}$$

Then,

$$\xi_\beta(x) \geq c \log(d(0,x)+1) - 4\beta S^2 JK \sum_{\ell=0}^{d(0,x)-1} (\cosh(c \log \frac{\ell+2}{\ell+1}) - 1)(\ell+1). \tag{5.3}$$

From Taylor expansions of the logarithm and of the hyperbolic cosine, there exist C, C' such that

$$\begin{aligned} \xi_\beta(x) &\geq c \log(d(0,x)+1) - 4\beta S^2 JK c^2 \sum_{\ell=1}^{d(0,x)} \frac{1}{\ell} - C' \\ &\geq [c - 4\beta S^2 JK c^2] \log(d(0,x)+1) - C. \end{aligned} \tag{5.4}$$

The optimal choice is $c = (8\beta S^2 JK)^{-1}$. □

Theorem 5.2. Assume that $J_{xy}^1 = J_{xy}^2$ for all $x, y \in \Lambda$. Then, for $i = 1, 2$, we have

$$|\langle S_0^i S_x^i \rangle| \leq S^2 e^{-\xi_\beta(x)}.$$

In the case of 2D-like graphs, we can use Lemma 5.1 and we obtain algebraic decay with a power greater than $(8\beta JS^2 K)^{-1}$.

Proof. We use the method of complex rotations. Let

$$S_y^\pm = S_y^1 \pm iS_y^2. \tag{5.5}$$

One can check that for any $a \in \mathbb{C}$, we have

$$e^{aS_y^3} S_y^\pm e^{-aS_y^3} = e^{\pm a} S_y^\pm. \tag{5.6}$$

We have $\langle S_0^+ S_x^- \rangle = 2\langle S_0^1 S_x^1 \rangle$ and this is non-negative by Theorem 4.1. Hamiltonian (4.1) can be rewritten as

$$H_\Lambda = -\frac{1}{2} \sum_{y,z \in \Lambda} (J_{yz}^1 S_y^+ S_z^- + J_{yz}^3 S_y^3 S_z^3). \tag{5.7}$$

Given numbers ϕ_y , let

$$A = \prod_{y \in \Lambda} e^{\phi_y S_y^3}. \tag{5.8}$$

Then,

$$\text{Tr } S_0^+ S_x^- e^{-\beta H_\Lambda} = \text{Tr } A S_0^+ S_x^- A^{-1} e^{-\beta A H_\Lambda A^{-1}}. \tag{5.9}$$

We now compute the rotated Hamiltonian,

$$\begin{aligned} A H_\Lambda A^{-1} &= -\frac{1}{2} \sum_{y,z \in \Lambda} (J_{yz}^1 e^{\phi_y - \phi_z} S_y^+ S_z^- + J_{yz}^3 S_y^3 S_z^3) \\ &= H_\Lambda - \frac{1}{2} \sum_{y,z \in \Lambda} J_{yz}^1 (\cosh(\phi_y - \phi_z) - 1) S_y^+ S_z^- - \frac{1}{2} \sum_{y,z \in \Lambda} J_{yz}^1 \sinh(\phi_y - \phi_z) S_y^+ S_z^- \\ &\equiv H_\Lambda + B + C. \end{aligned} \tag{5.10}$$

Notice that $B^* = B$ and $C^* = -C$. We obtain

$$\text{Tr } S_0^+ S_x^- e^{-\beta H_\Lambda} = e^{\phi_0 - \phi_x} \text{Tr } S_0^+ S_x^- e^{-\beta H_\Lambda - \beta B - \beta C}. \tag{5.11}$$

We now estimate the trace in the right side using the Trotter product formula and the Hölder inequality for traces. Recall that $\|B\|_s = (\text{Tr} |B|^s)^{1/s}$, with $\|B\|_\infty = \|B\|$ being the usual operator norm,

$$\begin{aligned} \text{Tr} S_0^+ S_x^- e^{-\beta H_\Lambda - \beta B - \beta C} &= \lim_{N \rightarrow \infty} \text{Tr} S_0^+ S_x^- \left(e^{-\frac{\beta}{N} H_\Lambda} e^{-\frac{\beta}{N} B} e^{-\frac{\beta}{N} C} \right)^N \\ &\leq \lim_{N \rightarrow \infty} \|S_0^+ S_x^-\|_\infty \|e^{-\frac{\beta}{N} H_\Lambda}\|_N^N \|e^{-\frac{\beta}{N} B}\|_\infty^N \|e^{-\frac{\beta}{N} C}\|_\infty^N. \end{aligned} \quad (5.12)$$

Observe now that $\|S_0^+ S_x^-\| = 2S^2$, $\|e^{-\frac{\beta}{N} H_\Lambda}\|_N^N = Z_\Lambda$, $\|e^{-\frac{\beta}{N} B}\|_\infty^N \leq e^{\beta \|B\|}$, and $\|e^{-\frac{\beta}{N} C}\|_\infty^N = 1$. The theorem then follows from

$$\|B\| \leq S^2 \sum_{y,z \in \Lambda} |J_{yz}^1| (\cosh(\phi_y - \phi_z) - 1). \quad (5.13)$$

□

ACKNOWLEDGMENTS

We are grateful to the Institute for Advanced Study at Princeton for hospitality. J.F. thanks Thomas Spencer for his generosity and interest and the “Fund for Math” and “The Robert and Luisa Fernholz Visiting Professorship Fund” for financial support.

- ¹ C. A. Bonato, J. Fernando Perez, and A. Klein, “The Mermin-Wagner phenomenon and cluster properties of one- and two-dimensional systems,” *J. Stat. Phys.* **29**, 159–175 (1982).
- ² C. Borgs, R. Kotecký, and D. Ueltschi, “Low temperature phase diagrams for quantum perturbations of classical spin systems,” *Commun. Math. Phys.* **181**, 409–446 (1996).
- ³ O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2* (Springer, 1981).
- ⁴ N. Datta, R. Fernández, and J. Fröhlich, “Low-temperature phase diagrams of quantum lattice systems. I. Stability for quantum perturbations of classical systems with finitely many ground states,” *J. Stat. Phys.* **84**, 455–534 (1996).
- ⁵ S. Dirren, Diploma thesis, ETH Zürich, 1999.
- ⁶ F. Dyson, E. H. Lieb, and B. Simon, “Phase transitions in quantum spin systems with isotropic and non isotropic interactions,” *J. Stat. Phys.* **18**, 335–383 (1978).
- ⁷ M. E. Fisher and D. Jasnow, “Decay of order in isotropic systems of restricted dimensionality. II. Spin systems,” *Phys. Rev. B* **3**, 907–924 (1971).
- ⁸ J. Fröhlich, R. Israel, E. H. Lieb, and B. Simon, “Phase transitions and reflection positivity. I. General theory and long range lattice models,” *Commun. Math. Phys.* **62**, 1–34 (1978).
- ⁹ J. Fröhlich and C.-É. Pfister, “On the absence of spontaneous symmetry breaking and of crystalline ordering in two-dimensional systems,” *Commun. Math. Phys.* **81**, 277–298 (1981).
- ¹⁰ J. Fröhlich and C.-É. Pfister, “Absence of crystalline ordering in two dimensions,” *Commun. Math. Phys.* **104**, 697–700 (1986).
- ¹¹ J. Fröhlich and P.-F. Rodriguez, “Some applications of the Lee-Yang theorem,” *J. Math. Phys.* **53**, 095218 (2012).
- ¹² C. Gruber and H. Kunz, “General properties of polymer systems,” *Commun. Math. Phys.* **22**, 133–161 (1971).
- ¹³ F. Guerra, L. Rosen, and B. Simon, “Correlation inequalities and the mass gap in $P(\phi)_2$,” *Commun. Math. Phys.* **41**, 19–32 (1975).
- ¹⁴ K. R. Ito, “Clustering in low-dimensional $SO(N)$ -invariant statistical models with long-range interactions,” *J. Stat. Phys.* **29**, 747–760 (1982).
- ¹⁵ T. Koma and H. Tasaki, “Decay of superconducting and magnetic correlations in one- and two-dimensional Hubbard models,” *Phys. Rev. Lett.* **68**, 3248–3251 (1992).
- ¹⁶ R. Kotecký and D. Preiss, “Cluster expansion for abstract polymer models,” *Commun. Math. Phys.* **103**, 491–498 (1986).
- ¹⁷ K. Kubo, “Existence of long-range order in the XY model,” *Phys. Rev. Lett.* **61**, 110–112 (1988).
- ¹⁸ J. L. Lebowitz and O. Penrose, “On the exponential decay of correlations,” *Commun. Math. Phys.* **39**, 165–184 (1974).
- ¹⁹ O. A. McBryan and T. Spencer, “On the decay of correlations in $SO(n)$ -symmetric ferromagnets,” *Commun. Math. Phys.* **53**, 299–302 (1977).
- ²⁰ C.-É. Pfister, “Analyticity properties of the correlation functions for the anisotropic Heisenberg model,” *Commun. Math. Phys.* **41**, 109–117 (1975).
- ²¹ B. Simon, *The Statistical Mechanics of Lattice Gases* (Princeton University Press, 1993).
- ²² D. Ueltschi, “Cluster expansions and correlation functions,” *Moscow Math. J.* **4**, 511–522 (2004).
- ²³ D. Ueltschi, “Random loop representations for quantum spin systems,” *J. Math. Phys.* **54**, 083301 (2013).